

**Rec'd PCT/PTO 02 0 MAR 2002**

FORM PTO-1390 (Modified) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (REV 11-2000)		ATTORNEY'S DOCKET NUMBER <b>220303US0XPCT</b>
<b>TRANSMITTAL LETTER TO THE UNITED STATES</b> <b>DESIGNATED/ELECTED OFFICE (DO/EO/US)</b> <b>CONCERNING A FILING UNDER 35 U.S.C. 371</b>		U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR <b>10/088072</b>
INTERNATIONAL APPLICATION NO. <b>PCT/CA00/01142</b>	INTERNATIONAL FILING DATE <b>3 October 2000</b>	PRIORITY DATE CLAIMED <b>5 October 1999</b>
TITLE OF INVENTION <b>RHODAMINE DERIVATIVES FOR PHOTODYNAMIC DIAGNOSIS AND TREATMENT</b>		
APPLICANT(S) FOR DO/EO/US <b>ROY Denis-Claude et al.</b>		
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: <ol style="list-style-type: none"> <li>1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below</li> <li>4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31).</li> <li>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c) (2))             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).</li> <li>b. <input checked="" type="checkbox"/> has been communicated by the International Bureau.</li> <li>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</li> </ol> </li> <li>6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> is attached hereto.</li> <li>b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4).</li> </ol> </li> <li>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</li> <li>b. <input type="checkbox"/> have been communicated by the International Bureau</li> <li>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</li> <li>d. <input checked="" type="checkbox"/> have not been made and will not be made.</li> </ol> </li> <li>8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</li> <li>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).</li> <li>10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).</li> <li>11. <input checked="" type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409).</li> <li>12. <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210).</li> </ol> <p><b>Items 13 to 20 below concern document(s) or information included:</b></p> <ol style="list-style-type: none"> <li>13. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</li> <li>14. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</li> <li>15. <input type="checkbox"/> A <b>FIRST</b> preliminary amendment.</li> <li>16. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</li> <li>17. <input type="checkbox"/> A substitute specification.</li> <li>18. <input type="checkbox"/> A change of power of attorney and/or address letter.</li> <li>19. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.</li> <li>20. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).</li> <li>21. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).</li> <li>22. <input type="checkbox"/> Certificate of Mailing by Express Mail</li> <li>23. <input checked="" type="checkbox"/> Other items or information:</li> </ol> <p style="margin-top: 10px;"> <b>Form PTO-1449</b>  <b>Request for Priority Under 35 U.S.C. 119(e)</b> </p>		

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.101) <div style="font-size: 1.5em; font-weight: bold;">10/088072</div>		INTERNATIONAL APPLICATION NO. <div style="font-weight: bold;">PCT/CA00/01142</div>		ATTORNEY'S DOCKET NUMBER <div style="font-weight: bold;">220303US0XPCT</div>	
--	--	---	--	---	--

24. The following fees are submitted: <b>BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :</b> <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... \$1040.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... \$890.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$740.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$100.00 <div style="text-align: right; font-weight: bold;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				<b>CALCULATIONS PTO USE ONLY</b>	
				\$890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).				\$130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	8 - 20 =	0	x \$18.00	\$0.00	
Independent claims	1 - 3 =	0	x \$84.00	\$0.00	
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>				\$0.00	
TOTAL OF ABOVE CALCULATIONS =				\$1,020.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$0.00	
SUBTOTAL =				\$1,020.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				\$0.00	
TOTAL NATIONAL FEE =				\$1,020.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00	
TOTAL FEES ENCLOSED =				\$1,020.00	
				Amount to be: refunded	\$
				charged	\$

a. ☒ A check in the amount of \$1,020.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.


c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 15-0030. A duplicate copy of this sheet is enclosed.

d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Surinder Sachar  
Registration No. 34,423

22850

Surinder Sachar

SIGNATURE \_\_\_\_\_

**Norman F. Oblon**

NAME \_\_\_\_\_

**24,618**

REGISTRATION NUMBER \_\_\_\_\_

March 20 2002

DATE \_\_\_\_\_

90 Rec'd PCT/PTO 3 JUL 2002

## APPLICATION DATA SHEET

## APPLICATION INFORMATION

Application Number:: 10/088,072  
Application Date:: 03/20/02  
Application Type:: REGULAR  
Subject Matter:: UTILITY  
CD-ROM or CD-R?: NONE  
Title:: RHODAMINE DERIVATIVES FOR  
PHOTODYNAMIC DIAGNOSIS AND  
TREATMENT  
Attorney Docket Number:: 220303US0XPCT

## INVENTOR INFORMATION

100  
Applicant Authority Type:: INVENTOR  
Primary Citizenship Country:: Canada ✓  
Status:: FULL CAPACITY  
Given Name:: Denis-Claude  
Family Name:: Roy  
City of Residence:: Laval  
State or Province of Residence:: Quebec  
Country of Residence:: Canada  
Street of Mailing Address:: 2444 Prudential  
City of Mailing Address:: Laval  
State or Province of Mailing Address:: Quebec  
Country of Mailing Address:: Canada  
Postal or Zip Code of Mailing Address:: H7K 2C4  
  
200  
Applicant Authority Type:: INVENTOR  
Primary Citizenship Country:: Canada ✓  
Status:: FULL CAPACITY  
Given Name:: Martin  
Family Name:: Guimond  
City of Residence:: Columbus  
State or Province of Residence:: Ohio  
Country of Residence:: United States  
Street of Mailing Address:: 1696 Quarry Trace  
City of Mailing Address:: Columbus  
State or Province of Mailing Address:: Ohio  
Country of Mailing Address:: United States  
Postal or Zip Code of Mailing Address:: 43204

388

Applicant Authority Type:: INVENTOR  
 Primary Citizenship Country:: Canada ✓  
 Status:: FULL CAPACITY  
 Given Name:: Nestor  
 Family Name:: Molfino  
 City of Residence:: Potomac  
 State or Province of Residence:: Maryland  
 Country of Residence:: United States  
 Street of Mailing Address:: 8817 Bells Mill Road  
 City of Mailing Address:: Potomac  
 State or Province of Mailing Address:: Maryland  
 Country of Mailing Address:: United States  
 Postal or Zip Code of Mailing Address:: 20854

## CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 22850

## REPRESENTATIVE INFORMATION

Representative Customer Number:: 22850

## DOMESTIC PRIORITY INFORMATION

Application::	Continuity Type::	Parent Application::	Parent Filing Date::
<u>This Application</u>	<u>National Stage of</u>	<u>PCT/CA00/01142</u>	<u>10/03/00</u>
<u>PCT/CA00/01142</u>	<u>Non-Provisional of</u>	<u>60/157,790</u>	<u>10/05/99</u>

## ASSIGNMENT INFORMATION

Assignee Name:: Universite de Montreal  
 Street of Mailing Address:: 2900 Edouard-Montpetit  
 City of Mailing Address:: Montreal  
 State or Province of Mailing Address:: Quebec  
 Country of Mailing Address:: Canada  
 Postal or Zip Code of Mailing Address:: H3T 1J4

Assignee Name:: Hopital Maisonneuve-Rosemont  
 Street of Mailing Address:: 5415, boul. l'Assomption  
 City of Mailing Address:: Montreal  
 State or Province of Mailing Address:: Quebec  
 Country of Mailing Address:: Canada  
 Postal or Zip Code of Mailing Address:: H1T 2M4

RHODAMINE DERIVATIVES FOR PHOTODYNAMIC DIAGNOSIS  
AND TREATMENT

BACKGROUND OF THE INVENTION

5 (a) Field of the Invention

The invention relates to a photodynamic treatment for the preferential destruction of immunologically reactive cells without substantially affecting the normal cells or causing systemic toxicity  
10 for the patient.

(b) Description of Prior Art

Immunologic disorders are conditions or diseases that result from the production of immune cells recognizing normal cells and tissues as foreign. Cells  
15 with immunoreactivity towards normal cells or tissues induce damages in these normal cells and tissues either directly, through cellular effector mechanisms, or indirectly through antibodies, cytokines or other  
20 mediators. Such immunologic disorders are usually divided in alloimmune conditions and autoimmune conditions. Alloimmune disorders occur primarily in the context of allogeneic transplantation (bone marrow and other organs: kidney, heart, liver, lung, etc.).  
25 In the setting of bone marrow transplantation, donor immune cells present in the hematopoietic stem cell graft react towards host normal tissues, causing graft-versus-host disease (GVHD). The GVHD induces damage primarily to the liver, skin, intestine, lung, eyes and  
30 mouth. Autoimmune disorders are comprised of a number of arthritic conditions, such as rheumatoid arthritis, scleroderma and lupus erythematosus; endocrine conditions, such as diabetes mellitus; neurologic conditions, such as multiple sclerosis and myasthenia  
35 gravis; gastrointestinal conditions, such as Crohn's disease and ulcerative colitis; hematological

disorders, such as autoimmune hemolytic anemia, etc. The immune reaction in both alloimmune and autoimmune disorders progresses to generate organ dysfunction and damage.

5 Despite important advances in treatment, immunologic complications remain the primary cause of failure of allogeneic transplantations, whether in hematopoietic stem cell transplantation (GVHD) or in solid organ transplantation (graft rejection). In addition, autoimmune disorders represent a major cause of both morbidity and mortality. Prevention and treatment of these immune disorders has relied mainly on the use of immunosuppressive agents, monoclonal antibody-based therapies, radiation therapy, and more recently molecular inhibitors. Significant improvement in outcome has occurred with the continued development of combined modalities, but for a small number of disorders and patients. However, for the most frequent types of transplantation (bone marrow, kidney, liver, heart and lung), and for most immune disorders (rheumatoid arthritis, connective tissue diseases, multiple sclerosis, etc.) resolution of the immunologic dysfunction and cure has not been achieved. Therefore, the development of new approaches for the prevention and treatment of immunologic disorders is critically needed particularly for those patients who are at high risk or whose disease has progressed and are refractory to standard immunosuppressive therapy. Allogeneic stem cell transplantation (AlloSCT) has been employed for the treatment of a number of malignant and non-malignant conditions. Allogeneic stem cell transplantation is based on the administration of high-dose chemotherapy with or without total body irradiation to eliminate malignant cells, and host hematopoietic cells. Normal hematopoietic donor stem

cells are then infused into the patient in order to replace the host hematopoietic system. AlloSCT has been shown to induce increased response rates when compared with standard therapeutic options. One  
5 important issue that needs to be stressed when using AlloSCT relates to the risk of reinfusing immune cells that will subsequently recognize patient cells as foreign and cause GVHD. A variety of techniques have been developed that can deplete up to 99,999% of T  
10 cells from the stem cell graft. These techniques, including immunologic and physical purging, are not entirely satisfactory. One major consideration when purging stem cell grafts is to preserve the non-host-reactive T cells so that they can exert anti-infectious  
15 and anti-leukemia activity upon grafting. The potential of photodynamic therapy, in association with photosensitizing molecules capable of destroying immunologically reactive cells while sparing normal donor-non-reactive immune cells to purge hematopoietic  
20 cell grafts in the preparation of AlloSCT or autologous stem cell transplantation (AutoSCT) and after AlloSCT in the context of donor lymphocyte infusions to eliminate recurring leukemia cells has largely been unexplored. To achieve eradication of T cells, several  
25 approaches have been proposed including:

- 1) *in vitro* exposure of the graft to monoclonal antibodies and immunotoxins against antigens present on the surface of T cells (anti-CD3, anti-CD6, anti-CD8, etc.);
- 30 2) *in vitro* selection by soybean agglutinin and sheep red blood cell rosetting;
- 3) positive selection of CD34+ stem cells with or without additional negative selection of T cells;

- 4) *in vivo* therapy with combinations of anti-thymocyte globulin, or monoclonal antibodies,
- 5) *in vivo* or *ex vivo* treatment with photosensitizing agents; and
- 5 6) *In vitro* or *ex vivo* exposure of recipient-reactive donor T cells by monoclonal antibodies or immunotoxins targeting the interleukin 2 receptor or OX-40 antigen (Cavazzana-Calvo M. et al. (1990) Transplantation, 50:1-7; Tittle T.V. et al (1997) Blood 89:4652-58; Harris D.T. et al. (1999) Bone Marrow Transplantation 23:137-44).
- 10

However, most of these methods are not specifically directed at the alloreactive T cell subset, but rather aiming at the elimination of either  
15 all T cells or broad T cell populations. This is associated with numerous problems, including disease recurrence, graft rejection, second malignancies and severe infections. In addition, the clinical relevance  
20 of several of these methods remains to be established.

There are many reports on the use of photodynamic therapy in the treatment of malignancies (Daniell M. D., Hill J. S. (1991) Aust. N. Z. J. Surg., 61: 340-348). One of these uses is described in U.S.  
25 Patents numbers 5,556,992 and 5,773,460, where novel photoactivable rhodamine derivatives are used for the photodynamic therapy of a cancer patient by destroying human cancer cells, wherein appropriate intracellular levels of the derivatives are achieved and irradiation  
30 with light of a suitable wavelength is applied. The method has been applied for cancers of various origins and for the eradication of viruses and pathogens (Raab O. (1990) Infusoria Z. Biol., 39: 524).

The initial experiments on the use of  
35 photodynamic therapy for cancer treatment using various



There is currently a lack of agents, which allow selective destruction of immunologic cells while leaving intact the normal non-pathogenic residual cellular population. Preferential uptake of photosensitive dye and cytotoxicity of photodynamic therapy against lymphoid cells (Greinix H.T., et al. Blood (1998) 92:3098-3104; Hunt D.W. et al (1999) Immunopharmacology, 41:31-44; Heykorenko E.A et al (1998) Immunopharmacology 40: 231-40); and macrophages

5

10

- 15

## 20

30

35

is taken up and is preferentially retained by many tumor cell types, impairing their proliferation and survival by altering membrane and mitochondrial function (Oseroff A. R. (1992) *In Photodynamic therapy* (Henderson B. W., Dougherty T. J. , eds) New York: Marcel Dekker, pp. 79-91). *In vivo*, chemotherapy with rhodamine 123 can prolong the survival of cancerous mice, but, despite initial attempts to utilize rhodamine 123 in the treatment of tumors, the systemic toxicity may limit its usefulness (Bernal, S.D., et al. (1983) *Science*, **222**: 169; Powers, S.K. et al. (1987) *J. Neurosur.*, **67**: 889).

United States Patent No. 4,612,007 issued on September 16, 1986 in the name of Richard L. Edelson, discloses a method for externally treating human blood, with the objective of reducing the functioning lymphocyte population in the blood system of a human subject. The blood, withdrawn from the subject, is passed through an ultraviolet radiation field in the presence of a dissolved photoactive agent capable of forming photoadducts with lymphocytic-DNA. This method presents the following disadvantages and deficiencies. The procedure described is based on the utilization of known commercially available photoactive chemical agents for externally treating patient's blood, leaving immune cells from other sites intact in the process. According to Richard L. Edelson, the method only reduces, does not eradicate, the target cell population. This treatment strategy does not incorporate any attempt to enhance the immunoreactivity of target cells. Moreover, the wavelength range of UV radiation used in the process proposed by Richard L. Edelson could be damageable to the normal cells.

International Application published on January 7, 1993 under International publication number WO

It would be highly desirable to be provided with a new use of rhodamine derivatives in the treatment of immunologic cells, which overcomes these drawbacks while having no substantial systemic toxicity for the patient.

**SUMMARY OF THE INVENTION**

One aim of the present invention is to provide new use of photosensitizers endowed with the following characteristics:

- i) preferential localization and uptake by the immunologic cells;
- ii) upon application of appropriate light intensities, functional or physical elimination of those cells which have accumulated and retained the photosensitizing agents;
- iii) sparing a sufficient proportion of the normal hematopoietic T and stem cell compartment from the destructive effects of activated photosensitizers;
- iv) utilization of photosensitizers for hematopoietic stem cell purging of immunologic cells in preparation for allogeneic or autologous stem cell transplantation with or without the use of strategies to increase immunoreactivity; and
- v) utilization of photosensitizers for ex vivo elimination of reactive immune cells in patients with immunological disorders with or without the use of strategies.
- vi) utilization of photosensitizers to evaluate transport mechanism of immune and malignant cells.

In accordance with the present invention, there is provided a photoactivable pharmaceutical composition for the selective destruction and/or inactivation of immunologically reactive cells without substantially affecting the normal cells or causing systemic toxicity for the patient, the composition comprising at least

30 In accordance with the present invention, there is provided a method of prevention of graft-versus-host disease associated with allogeneic stem cell transplantation in a patient, which comprises the steps of:

The method in accordance with a preferred embodiment of the present invention, wherein the  
35 hematopoietic cells is selected from the group

Fig. 3 demonstrates that cells from subject A  
35 activated against subject B cells and photodynamically



treated, do not proliferate when reexposed to B cells but do proliferate when exposed to C cells. A, B and C cells were from unrelated individuals: A and B differed by 2 HLA antigens (B and DR).

5        Fig. 4 shows TH9402 fluorescence upon flow cytometric evaluation of resting and activated lymphocytes. Cells were evaluated at various times after the end of the TH9402 incorporation period. Activated lymphocytes retain more TH9402 than resting  
10    lymphocytes.

      Fig. 5 shows the impact of cyclosporin A on the TH9402 cellular efflux after 110 minutes from the end of the TH9402 incorporation period. Cyclosporin A blocks the efflux of TH9402 in resting lymphocytes, but  
15    not in activated lymphocytes.

      Fig. 6 shows the effect of PDT with TH9402 on CD4 and CD8 positive cells after activation in mixed lymphocyte culture with third party cells. Activated cells (expressing CD25), both CD4+ and CD8+, are  
20    eliminated by photodynamic therapy.

      Figs. 7A and 7B show that approximately 3 logarithms (99.9%) of human B cells can be eliminated by PDT with TF9402 (A). In contrast, less than one logarithm (approximately 50%) of hematopoietic  
25    progenitor cells of myeloid (colony forming units-granulocyte monocyte [CFU-GM], erythroid (burst forming units-erythroid [BFU-E], and mixed (colony forming units-granulocyte erythrocyte monocyte megakaryocyte [CFU-GEMM]) origin are eliminated by the same PDT  
30    procedure.

      Figs. 8A, 8B, and 8C show three graphs of the photo toxicity of 4,5-dibromorhodamine 110 n-butyl ester hydrobromide salt used in accordance with the method of the present invention and expressed in %  
35    viability.

Figs. 9A and 9B show two graphs of the photo toxicity of rhodamine B n-butyl ester hydrochloride salt used in accordance with the method of the present invention and expressed in % viability.

5

#### DETAILED DESCRIPTION OF THE INVENTION

Photoactive dyes are excited from the ground state to the singlet excited state following absorption of photons. Singlet excited states of organic molecules generally have short lifetimes ( $10^{-12}$ - $10^{-6}$  sec.) as they rapidly relax back to the ground state using non-radiative (vibrational modes) and radiative (fluorescence) processes. Intersystem crossing to the more stable triplet excited state is also competing with relaxation to the ground state. Triplet excited states generally have longer lifetimes ( $10^{-6}$ - $10$  sec) which allow them to diffuse and react with other molecules in the medium.

Triplet excited states can react with molecular oxygen via two different mechanisms. The first mechanism (Type I) consists of the transfer of an electron from the excited dyes to molecular oxygen, resulting in highly reactive free radical-anions being present in the cellular environment.

The second mechanism (Type II) consists of the transfer of energy from the excited dyes to molecular oxygen, leading to the formation of cytotoxic singlet oxygen.

Photosensitizers must therefore meet two conditions in order to be effective phototherapeutic agents. The first condition is that they must be present at a higher concentration in target cells than in normal cells. A higher concentration of dyes in malignant and immunologic cells results in a higher amount of photogenerated cytotoxic species and

35



Different heavy atom substitutions of the hydrogen atoms (halogenic substitution) of the rhodamine backbone, for example, dibromo and diiodo derivatives of rhodamine B and rhodamine 110 esters, were prepared and tested.

Substitution of the oxygen heteroatom of the rhodamine backbone by a heavier atom to reduce  $S_0/S_1$  splitting, theoretically should increase spin orbit coupling and promote intersystem crossing from the  $S_1$  to the  $T_1$  state, producing higher triplet yields than the original dye. This should increase proportionally the production of singlet oxygen. Therefore, S (Sulfur), Se (Selenium) and Te (Tellurium) substitutions for the oxygen atom (O) of the rhodamine backbone is being explored.

In accordance with the present invention, there is also shown that TH9402 is preferentially retained by activated T cells. Resting T cells can eliminate TH9402 from their intracellular milieu, but not activated T

cells (Fig.4). In addition, we found that TH9402 efflux is inhibited when cyclosporin-A is added (Fig. 5). Since cyclosporin-A is a potent inhibitor of Pgp-170, it is likely that TH9402 efflux relies on a Pgp-170 transporter, as previously observed for the rhodamine parent molecule. Inactivation of the MDR pathway in activated T cells could therefore explain the preferential elimination of activated T cells and preservation of unactivated T cells for subsequent recognition of third party cells (Fig.3). The absence of known strong expression of Pgp-170 on B cells prompted us to evaluate the capacity of PDT with TH9402 to eliminate B lymphocytes. TH9402 was indeed found capable of eliminating approximately 3 logarithms (99.9%) of B lymphocytes. In contrast, more than half of normal hematopoietic progenitors of myeloid (CFU-GM), erythroid (BFU-E) and mixed (CFU-GEMM) origin are preserved when PDT is performed in the same conditions as used to obtain high levels of elimination of B lymphocytes. Therefore, PDT with TH9402 presents a therapeutic profile favorable to the elimination of immune cells, including activated T cells, B cells and potentially other cells (such as dendritic cells) that could be involved in immune disorders. The photodynamic treatment herein described could be done in conjunction with prior sensitization or activation of potential effector cells, or without manipulations to increase immunoreactivity since pathogenic immune cells may (1) be already activated because of the underlying disease, or (2) be spontaneously sensitive to PDT (e.g. B cells). Activation could be achieved through exposure to antigens, cells, cell lysates, proteins, peptides, DNA, cytokines, mitogens, lectins, or other directly or indirectly activating processes.

In accordance with the present invention, there is provided the use of such above-mentioned dyes in conjugation with antibodies specific for immune cell populations, peptides, proteins, or poisonous  
5 substances, or liposomal or lipoproteins, inhibitors of efflux pathways (e.g MDR) or fluorochrome adducts or other agents.

In addition, the photosensitizers to be described have the potential to act synergistically in  
10 conjunction with other photoactive substances.

Moreover, the negative selection procedure provided by the use of photodynamic treatment does not preclude the use of other means for enriching hematopoietic stem cells such as positive selection  
15 with anti-CD34 monoclonal antibodies.

### **Clinical applications**

The first clinical application of the current invention is the use of photosensitizers in the context  
20 of in vitro purging of alloreactive cells prior to allogeneic stem cell transplantation for the prevention of graft-versus-host disease. In this condition, donor cells are first exposed to recipient cells or antigens or other components, in order to activate donor cells  
25 against antigens of the recipient. These cells then undergo photodynamic therapy to eliminate alloreactive donor cells. This strategy preserves hematopoietic cells that are non-reactive against host cells.

The same strategy (elimination of alloreactive  
30 cells from cellular grafts) could be applied in all instances where the administration of donor cells could induce graft-versus-host disease, such as in cases where donor lymphocytes are infused into recipients to exert anti-leukemia or anti-infection activity.

35 In the case of autoimmune disorders, a portion of immune cells are autoreactive. When autologous stem



chemotherapy, photodynamic or other therapeutic agents).

### Chemical Synthesis

5           The chemical synthesis of rhodamine B n-buthylester hydrochloride, 4,5-dibromorhodamine n-butylester hydrobromide, rhodamine n-buthylester hydrochloride, 4,5-dibromorhodamine 110 n-butulester hydrobromide and 4,5-dibromorhodamine 123 hydrobromide  
10 was effected as described in US Patent No. 5,556,992 issued on September 17, 1996, which is hereby incorporated by reference.

### Cell lines

15           T cells represent the most important population of immune cells present in the peripheral blood. In order to demonstrate the efficacy of photodynamic therapy with TH9402 to eliminate activated T cells, we first evaluated its effect on a malignant T cell line.  
20 Phototoxicity was also evaluated in parallel against the chronic myelogenous leukemia cell line K562, that had been used in United States Patents Nos. 5,556,992 and 5,773,460. The CEM T cell acute lymphoblastic leukemia cell line and K562 chronic myelogeneous  
25 leukemia cell line (Lozzio,B.B. and Lozzio,C.B. (1979) *Cancer Res.*, 3(6): 363-370) were obtained from the American Type Culture Collection (ATCC, 12301 Parklawn Drive, Rockville, MD 20852 USA) under the accession number CCL-119 and CCL-243. Cultures were maintained  
30 at 37°C in a humidified incubator with an atmosphere of 95% air and 5% CO<sub>2</sub>. Cells were grown in RPMI 1640 medium (GIBCO, Grand Island, NY)) supplemented with 10% fetal bovine serum (FBS), 2mM L-glutamine, 1mM sodium pyruvate, 100U/mL penicillin, and 100µg/mL streptomycin  
35 (Life Technologies, Inc.). Before each experiment, cell viability was assessed by trypan blue exclusion. CEM



5

## Photodynamic treatment

10

15

20

25

Phototoxicity of 4,5-dibromorhodamine 123 hydrobromide  
(TH9402)

30

39

treatment, cells were washed 3 times and plated in a limiting dilution assay (LDA) as described previously (Roy DC et al, *JNCI* 1996;88:1136-45). Briefly, each treatment sample was serially diluted from  $5 \times 10^5$  to 0.5  
5 cells per 100  $\mu$ l in RPMI 1640 supplemented with 10% FBS. Then, 24 aliquots of each dilution were plated in flat bottom microculture plates (Nunc, Nunc, Denmark). Cells were fed every 4 days and incubated at 37°C for 12-14 days. Growth at each serial dilution  
10 was assessed in an "all-or-nothing" (positive or negative) fashion under an inverted phase microscope. Frequency of clonogenic cells within the test population was estimated using chi-square minimization (Taswell C, *J.Immunol.* 1981;126:1614-19). As shown in  
15 Fig.1, photodynamic therapy with TH9402 eliminated almost all CEM and K562 cells, with less than 0.1% of CEM and K562 escaping elimination by phototherapy in comparison to the media only sample. These results indicate high levels of elimination of malignant T  
20 cells, as was previously reported for leukemic K562 cells, and support efficacy of this procedure for the elimination of malignant T cells. TH9402 was shown to be highly phototoxic; the elevated level of cytotoxic activity is believed to be a consequence of increased  
25 intracellular content of TH9402 in these malignant T and myeloid cell lines.

#### **T cell activation with PHA.**

30 Normal PBMC were activated by incubation at 37°C for 48 to 72 hours in X-vivo-15 medium (Biowhittaker, Walkersville. Md. U.S.A.) supplemented with 20% AB serum (Sigma), 1% pen-strep (Gibco), 2% glutamin (Gibco) and 20  $\mu$ g/ml of phytohemagglutinin-A (PHA-P)  
35 (Sigma). Cells were cultured in 25cm<sup>2</sup> flasks at a concentration of  $3 \times 10^6$  cells/ml. Following incubation,

cells were washed and treated with the TH9402 photodynamic treatment as described above, and proliferative activity measured as described below.

5 **Proliferation assay (mixed lymphocyte reaction)**

To evaluate the residual proliferative potential of activated mononuclear cells after photodynamic therapy, peripheral blood mononuclear cells were placed in 96-well microtiter plates and were incubated with  
10 PBMC from various individuals (demonstrating at least 2 major histocompatibility complex antigen mismatches with treated cells). The latter cells were serially diluted in order to obtain effector (treated cells) to target ratios ranging from 2:1 to 1:4 ( $4 \times 10^4$  treated  
15 effector cells/well) and incubated at 37°C for 5 days. Eighteen hours prior to harvesting, 1  $\mu$ Ci of  $^3$ H-thymidine was added. Cells were harvested using a PHD cell harvester (Cambridge Technology, Boston, MA, USA). Radioactivity in the cell harvest was counted using a  
20 liquid scintillation counter (Beckman, Chicago, IL, USA).

**Phototoxicity of 4,5-dibromorhodamine 123 hydrobromide against PHA activated cells**

25 The phototoxicity of TH9402 against PHA-activated PB mononuclear cells was assessed after photodynamic treatment using 5 and 7.5 joules/cm<sup>2</sup> of light energy (Fig. 2). After treatment, the cells were washed and evaluated for proliferative activity in a  
30 mixed lymphocyte reaction, according to the protocol in the previous paragraph. In PHA-activated cells that did not receive photodynamic therapy (untreated), proliferation in mixed lymphocyte culture increased with the number of effector cells. In contrast, when  
35 PHA-activated cells were treated with TH9402 using 5 and 7.5 J/cm<sup>2</sup> light energies, reactivity towards MHC

incompatible cells was abrogated. This result indicates that photodynamic therapy of PHA-activated cells is a very potent inhibitor of immunoreactivity in these cells. Cell counts performed three days after  
5 the photodynamic treatment show a decrease by more than ninety percent (90%) of the treated cells in comparison to the medium control. These results indicate that the loss of proliferative activity in activated cells is most likely due to the elimination of effector cells.

10

#### Allogeneic T cell activation

Another approach was used in this study to activate cells against specific target antigens. Mononuclear cells from subject A were incubated with  
15 irradiated mononuclear cells from subject B. In this one-way mixed lymphocyte culture, subjects A and B were unrelated and showed only partial human leukocyte antigen (HLA) matching with differences at two major histocompatibility complex (MHC) antigens. Briefly,  
20  $25 \times 10^6$  PBMC were incubated at  $37^\circ\text{C}$  for 4 days with  $25 \times 10^6$  irradiated (25Gy) stimulating mononuclear cells in X-vivo-15 medium (BioWhittaker) supplemented with 20% AB serum (Sigma), 1% pen-strep (Gibco), 2% glutamin (Gibco) and 50U/ml of IL-2 (ID lab). All cultures were  
25 performed in  $75\text{cm}^2$  flasks (Corning) in a final volume of 25 ml. The unstimulated control was performed with  $25 \times 10^6$  irradiated autologous cells.

After this activation period, cells had photodynamic therapy with TH9402 as described above.  
30 Following treatment, cells were plated in a proliferation assay for 5 days as described above where targets consisted of PBMC from subject B and also from subject C (mismatched unrelated). As shown in Figure

3, when cells from subject A, activated against B, underwent TH9402 photodynamic therapy, they did not proliferate when reexposed to cells from B. However, when the same A cells were exposed to C cells, they had  
 5 retained the capacity to proliferate. These results indicate that photodynamic therapy can specifically eliminate alloreactive cells, while sparing the alloreactive potential of unactivated cells. In addition, they demonstrate that it is possible to take  
 10 advantage of this activation strategy to deplete immunologic populations against a desired antigen.

#### Cellular concentration of TH9402

TH9402 cellular content in resting and activated  
 15 lymphocytes was evaluated by flow-cytometry, since the intensity of TH9402 (green) fluorescence correlates with the cellular content in TH9402. Briefly,  $10^6$  cells/ml, previously activated or not with PHA, were incubated in X-vivo-15 medium supplemented with 2.5%  
 20 human AB serum and  $10\mu\text{M}$  TH9402 for 40 minutes. These cells were washed two times with X-vivo medium supplemented with 10% AB serum and cells analysed by flow cytometry 30, 50, 70, 90 and 110 min after the end of the TH9402 incorporation period. As shown in Figure  
 25 4, resting lymphocytes rapidly lost TH9402 with approximately 50% (fifty percent) of cells demonstrating low TH9402 fluorescence 110 minutes after the end of the incorporation period. In addition, at all time-points evaluated, the intensity of TH9402  
 30 fluorescence was less for resting lymphocytes than for activated lymphocytes (Fig. 4). Since cellular concentration of TH9402 correlates with the extent of

cell elimination, the high concentration of TH9402 maintained in activated lymphocytes explains their sensitivity to photodynamic therapy. In contrast, the rapid efflux of TH9402 from resting lymphocytes should  
5 explain preservation of their proliferative activity.

In order to identify the mechanism responsible for the differential retention of TH9402 between activated and resting lymphocytes, cyclosporin-A was used to block the multidrug transporter (P-gp 170).  
10 These cells were incubated with 10 $\mu$ M TH9402 for 8 minutes, and washed with medium containing 1 $\mu$ g/ml cyclosporin-A or medium alone. TH9402 retention was subsequently evaluated by flow-cytometry (green fluorescence) (Fig. 5). After 110 minutes from the end  
15 of TH9402 incorporation, fluorescence intensity was identical in activated cells treated or not treated with cyclosporin A. In contrast, cyclosporin A induced higher retention of TH9402 in resting lymphocytes, suggesting that a functional P-gp is involved in TH9402  
20 dye efflux from resting lymphocytes and represents a major mechanism whereby these cells escape elimination by photodynamic therapy. The functional impairment of such a pump in activated lymphocytes could explain the high levels of phototoxicity observed in these cells.

25 **Phenotypic analysis of residual T lymphocytes after phototherapy using TH9402**

In order to determine if the abrogation of reactivity toward subject B obtained after PDT correlated with the  
30 loss of activated T cells, the proportion of activated cells was determined in samples exposed or not to PDT. Activated cells can be discriminated from resting T lymphocytes by their enhanced expression of CD25, which can be detected with a monoclonal antibody specific for  
35 CD25, the inducible  $\alpha$  chain of the IL-2 receptor.

Differential phototoxic activity of TH9402 against B  
35 cells and non-lymphoid hematopoietic progenitors

To evaluate the potential of PDT with TH9402 to eliminate other immune cell populations, normal human B cells were used as targets. Mononuclear cells from  
 5 normal donors were obtained by leukopheresis, and resuspended at 20 million cells per ml during the whole PDT process. Cells were centrifuged and resuspended in pre-warmed (37°C) X-Vivo-15 medium supplemented with 2.5% FCS and 10 U/ml heparin, with 5µM TH9402. After 40  
 10 minutes of incubation at 37°C, cells were washed and resuspended in a X-Vivo-15 medium and 10% FCS with 10 U/ml heparin (medium free of TH9402) for an efflux period of 50 minutes before exposure to light energy (10 to 30 Joules/cm<sup>2</sup>). Cells underwent light exposure  
 15 at 20 million cells per ml and at a thickness of 2 cm.

To evaluate the capacity of the PDP treatment to eliminate B cells, we used an *in vitro* B cell culture system. Briefly, 5x10<sup>6</sup> untreated and treated mononuclear cells were added to a 25 mm<sup>2</sup> monolayer of  
 20 irradiated mouse fibroblasts NIH 3T3 transfected to express CD40 ligand, an important molecule for B cells activation and proliferation. The cells were cultured during seven days in interleukin-4 (IL-4) (100 u/ml) containing-medium (Iscove's Modified Dulbecco Medium-1  
 25 MDM) with 2% FCS, 1% penicillin-streptomycin, 50 µg/ml human transferrin, 0.5% BSA, 5 µg/ml bovine insulin, 50 µg/ml of each oleic, linoleic and palmitic acid). At the end of the culture period, a trypan blue viability test was done as well as an immunophenotypic analysis  
 30 of residual CD19+ cells by flow-cytometry as described above.

To verify that the treatment preserved normal hematopoietic progenitors, we have used a clonogenic assay to measure the amount of hematopoietic clonogenic  
 35 precursors present in the same samples. Briefly, after PDP, all samples, including controls, were diluted and



plated at various cells densities (10,000 to 800,000) cells in semi-solid methylcellulose medium (StemCell Technologies Inc). Colonies were enumerated for myeloid, erythroid, and mixed progenitors after 13 to  
 5 16 days of incubation at 37°C, 5% CO<sub>2</sub>, and 98% relative humidity. Assays were done at least in duplicate. To determine the comparative reduction of the precursor cells, the mean values for each PDP condition were converted to percent of the appropriate control.

10 Normal human mononuclear cells were obtained and subjected to various PDP conditions to determine the efficacy of eradication of B cells, specificity and safety of the procedure. The number of B cells eliminated by TH9402 PDT increased with the level of  
 15 light energy delivered (Fig.7A). In comparison to untreated cells, PDT resulted in approximately 3 logarithms (99.9%) of eradication of B cells. In contrast, when these cells were evaluated for the elimination of non-lymphoid hematopoietic progenitors,  
 20 usually less than 50% (half of a logarithm) of these progenitors were eliminated by the same conditions of PDT (Fig.7B). These results indicate that immune cells other than activated T cells, such as B cells, can be eliminated by PDT with TH9402. In addition,  
 25 preservation of a large proportion of CFU-GM, BFU-E and CFU-GEMM progenitors demonstrates the specificity of this PDT process for defined immune cell populations. In addition, it confirms the capacity of such PDT to preserve normal hematopoietic progenitor cells for  
 30 hematologic reconstitution when used in the context of purging of grafts prior to autologous or allogeneic transplantation.

35 **Phototoxicity of 4,5-dibromorhodamine 110 n-butyl ester hydrobromide**

To ascertain the photochemotherapeutic potential of 4,5-dibromorhodamine 110 n-butyl ester hydrobromide (DBBE), *in vitro* phototoxicity was evaluated in the K-562 cell line procedure described. The cells were  
5 incubated with increasing concentrations of DBBE and the cell viability was measured at different time points following photodynamic therapy. The results shown in Figs. 8A, 8B and 8C show that a dosage of 10  $\mu$ g/ml of the dye and a brief exposure to 514.5 nm  
10 radiation from an argon ion laser at 0.5 J/cm<sup>2</sup> completely suppress cell viability in less than 24 hours after irradiation.

#### 15 Phototoxicity of Rhodamine B n-butyl ester hydrochloride

The photo toxicity *in vitro* of rhodamine B n-butyl ester (RBBE) was evaluated in the K-562 cell line procedure, in order to assess its photochemotherapeutic potential. Comparison was made to the induced  
20 phototoxicity of rhodamine 123 (RH123) and of rhodamine nB-butyl ester hydrochloride. Cell viability was evaluated 2 and 20 hours after photodynamic therapy. The results shown in Figs. 9A and 9B demonstrate that a dosage of 10  $\mu$ g/ml of the dye and a photo exposure of 5  
25 J/cm<sup>2</sup> from argon ion laser (514.5 nm) significantly suppress cell viability of K562 cells in less than 20 hours after irradiation. Rhodamine 123 has no effect on cell viability, even at exposures of 5 J/cm<sup>2</sup>. Phototoxicity of 4,5-dibromorhodamine 110 n-butyl ester  
30 hydrobromide and rhodamine B n-butyl ester hydrochloride were only assessed against the cell line K562. However, we anticipate that their activity will be similar against T cells.

#### 35 Phototoxicity against hematopoietic progenitor cell cultures

Conventional approaches for the prevention and treatment of immunologic disorders such as immunosuppressive agents, radiotherapy and monoclonal antibody-based therapies are limited by their intrinsic toxicity and myelosuppressive effects. The introduction of strategies to eliminate T cells *in vitro* or *in vivo* has resulted in a decreased incidence of graft-versus-host disease after allogeneic stem cell transplantation, improved graft survival in solid organ transplantation and improved clinical conditions for patients with immunologic disorders. However, T cell depletion is associated with an increased incidence of infections and malignancies or recurrence of malignant diseases, which have limited the use of T cell

elimination strategies. These complications are primarily attributable to the non-specific elimination of a majority of T cells, which are responsible for the control of infection and anti-leukemia activity. To  
5 overcome these limitations and to expand the number of patients and age limit for intensive curative therapy, the potential benefit of selective *in vitro* elimination of immunologic cells prior to allogeneic stem cell transplantation has become widely acknowledged.  
10 Moreover, selective elimination of immunologic cells has the potential to be most useful in the context of donor lymphocyte infusion after transplantation, solid organ transplantation, and autoimmune disorders where the patient might benefit from the elimination of  
15 alloreactive or activated immune cell populations.

In an effort to develop new anti-neoplastic drugs that would allow selective destruction of alloreactive or activated immune cells, new dye molecules have been prepared and tested as possible new  
20 photosensitizers, useful for the photodynamic prevention and therapy of immunologic disorders. Three new photosensitizers of the pyrylium family were prepared and their cytotoxicity profile, which is similar to that of TH9402, provides evidence for their  
25 potential use in the photodynamic treatment of immunologic disorders and also in the prevention and/or treatment of graft-versus-host disease.

The present invention will be more readily understood by referring to the following examples which  
30 are given to illustrate the invention rather than to limit its scope.

## Method of prevention of graft-versus-host disease in the context of allogeneic stem cell transplantation

Allogeneic stem cell transplantation is performed for numerous neoplastic and non-neoplastic conditions. Hematological malignancies are comprised of leukemia, lymphoma, multiple myeloma, myelodysplastic syndromes, etc.; and non-hematological malignancies: aplastic anemia, congenital disorders, severe immunodeficiency syndromes, rheumatoid arthritis, scleroderma, lupus erythematosus, multiple sclerosis, HIV and other immune disorders.

GVHD develops against antigens present on recipient cells but not on donor cells. Immunological differences between donor and recipient could be present at the level of major histocompatibility antigens, minor histocompatibility antigens or tumor-associated antigens. Disparity is established using one or more of the following procedures on blood or bone marrow cells:

- a) HLA typing: conventional serologic typing or  
35 molecular to identify disparities between donor

and recipient in major histocompatibility complex class I and class II antigens; and

b) Mixed lymphocyte culture to identify differences in class II antigens; and

5 c) Minor histocompatibility antigens: although a few cytotoxic T cell lines are available and could be used to identify minor histocompatibility antigens, currently, these tests are only available for research purposes.

10

#### **Progenitor cell harvesting**

After diagnosis, bone marrow (BM) or peripheral blood (PB) or cord-blood derived hematopoietic stem cells from the donor is harvested using previously  
15 described procedures for allogeneic progenitor cell transplantation (reviewed in *Bone Marrow Transplantation*, Forman SJ, Blume KG, Thomas ED eds, Blackwell Scientific Publications, Cambridge MA, USA, 1994). Donor hematopoietic stem cells collected for  
20 allografting can be immediately incubated with irradiated (25Gy) host mononuclear or other cells. Host cells admixed with donor cells are incubated in sterile dye free medium supplemented with 20% autologous serum and interleukin-2 for 2 to 5 days.  
25 This procedure elicits donor cell alloreactivity towards the host, and the cell graft subsequently undergoes photodynamic treatment ex vivo as described below.

#### **30 Selective in vitro purging of immunologic cells**

Ex vivo treatment consist of short-term incubation of previously activated BM or PB stem cells with one or several of the selected photoactive compounds. Duration of incubation, cell concentration  
35 and drug molarity is determined for each patient using an aliquot of the harvested cell population. Excess of

dyes is removed by cell washes with sterile dye free medium supplemented with 2% autologous serum. Cells are next being exposed to radiant energy of sufficient intensities to effect photodynamic purging of immune  
5 cells. Efficacy of the photodynamic purging procedure is verified on an aliquot of the treated cell population, before cryopreservation and/or re-infusion to the patient is performed. Until re-infusion to the patient, the cells can be cryopreserved in 10%  
10 dimethylsulfoxide (DMSO) and 90% autologous serum, at -196°C in the vapor phase of liquid nitrogen.

#### **Systemic treatment of patients**

Following stem cell harvest, the patient is  
15 submitted to dose-intensive chemotherapy and/or irradiation when indicated.

#### **Allogeneic stem cell transplantation**

Following appropriate treatment of the patient  
20 by high-dose chemotherapy and/or irradiation and at the appropriate clinical moment, cryopreserved marrow or peripheral blood or cord blood stem cells will be rapidly thawed and returned to the patient.

25           *Ex vivo* treatment consists of short-term  
incubation of PB mononuclear cells with one or several  
of the selected photoactive compounds. Duration of  
incubation, cell concentration and drug molarity are  
determined for each patient using an aliquot of the  
30 harvested cell population. Excess of dyes is removed  
by cell washes in sterile dye free medium supplemented  
with 2% autologous serum. Cells are next being exposed  
to radiant energy of sufficient intensities to effect  
photodynamic purging of activated cells which mediate  
35 GVHD.



# **Administration of photodynamically treated cells to patients**

Leukopheresed cells that are photodynamically treated are reinfused into the patient. This approach enables the elimination of a large number of circulating activated lymphocytes and other cells involved in GVHD. In addition, cells spared by the photodynamic treatment are unactivated and their reinfusion into the patient may help restore normal immunologic equilibrium and induce immunomodulation.

## **EXAMPLE III**

### **Method of treatment of immunologic disorders**

#### **Diagnostic procedures**

Diagnosis of autoimmune disorders is established using conventional clinical, biochemical and/or histopathological examination of the blood or appropriate tissues. Severe autoimmune diseases are amenable to autologous transplantation (reviewed in Sullivan KM et al., *Am. Soc. Hematol., Educ. Program Book*, 1998:198-214).

#### **Harvesting of hematopoietic stem cells**

After diagnosis, bone marrow (BM), peripheral blood (PB) or cord blood (CB) mononuclear cells are harvested using previously described procedures for the autologous marrow transplantation in cancer therapy (reviewed in *Bone Marrow Transplantation*, Forman SJ, Blume KG, Thomas ED eds, Blackwell Scientific Publications, Cambridge MA, USA, 1994). Patient's hematopoietic stem cells collected for autograft are treated immediately ex vivo as described below.

#### **In vitro elimination of cells mediating autoimmune disorders**

*Ex vivo* treatment consists of short-term incubation of BM or PB stem cells with one or several of the selected photoactive compounds. Duration of incubation, cell concentration and drug molarity is  
5 determined for each patient using an aliquot of the harvested cell population. Excess of dyes is removed by cell washes in sterile dye free medium supplemented with 2% autologous serum. Cells are next being exposed to radiant energy of sufficient intensities to effect  
10 photodynamic purging of immunologic cells which mediate the immunologic disorder.

#### **Administration of photodynamically treated cells to patients**

15 Hematopoietic stem cells that are photodynamically treated are stored (frozen or kept in culture). This approach enables the elimination of a large number of activated lymphocytes and other cells involved in the immunologic disorder. In addition,  
20 cells spared by the photodynamic treatment are unactivated and their reinfusion may help restore normal immunologic equilibrium. Following stem cell harvest, patient are either treated with conventional regimens until autografting is clinically indicated or  
25 immediately submitted to dose-intensive chemotherapy and total body irradiation where indicated.

#### **Autologous stem cell transplantation**

Following high-dose chemotherapy and irradiation  
30 cryopreserved marrow or peripheral blood stem cells are rapidly thawed and infused to the patient.



clinical studies. This is useful for the investigation of cellular and molecular biology. Because MDR and other similar transporters can limit the activity or various therapeutic agents such as chemotherapeutic and photodynamic agents, this test should have diagnostic and prognostic importance and help identify optimum therapeutic strategies for patients with immunologic and neoplastic disorders.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

NO. 9406 P. 10/17

**WE CLAIM:**

2. Use of the photoactivable derivatives of claim 1 for the photodynamic treatment for the selective destruction and/or inactivation of immunologically reactive cells without substantially affecting the normal cells or causing systemic toxicity for the patient, wherein appropriate intracellular levels of said derivatives are achieved and irradiation of a suitable wavelength and intensity is applied.

DEC. 20. 2001 3:44PM

SWABEY OGILVY MTL 514 288 8389

NO. 9406 P. 11/17

- 42 -

3. A method of prevention of graft-versus-host disease associated with allogeneic stem cell transplantation in a patient, which comprises the steps of;

- a) activating lymphocytes from a donor by mixing donor cells with host cells for a period of time sufficient for an immune reaction to occur;
- b) substantially eliminating the activated lymphocytes of step a) with photodynamic therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and
- c) performing allogenic stem cell transplantation using the treated mix of step b).

4. A method for the treatment of immunologic disorder in a patient, which comprises the steps of:

- a) harvesting said patient's hematopoietic cells;
- b) ex vivo treating of the hematopoietic cells of step a) by photodynamic therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and
- c) performing graft infusion or autograft transplantation using the treated hematopoietic cells of step b).

5. The method of claim 4, wherein said immunologic disorder is selected from the group consisting of conditions in which self cells or donor cells react against host tissues or foreign targets, such as graft-versus-host disease, graft rejection, autoimmune disorders and immunoallergic conditions.

20-12-2001

CA000114

DEC. 20. 2001 3:44PM SWABEY OGILVY MTL 514 288 8389

NO. 9406 P. 12/17

- 43 -

6. The method of claim 4, wherein said hematopoietic cells is selected from the group consisting of bone marrow, peripheral blood, and cord blood mononuclear cells.

7. A method of evaluating transport mechanism of immune and/or malignant cells, which comprises using the photoactivable pharmaceutical composition of claim 1.

8. The method of claim 7, wherein said composition is evaluated by flow cytometry.

9. Use of a composition according to claim 1 in the photodynamic treatment for the selective destruction and/or inactivation of immunologically reactive cells without substantially affecting the normal cells or causing systemic toxicity for the patient, wherein appropriate intracellular levels of said derivatives are achieved and irradiation of a suitable wavelength and intensity is applied.

10. Use of a composition according to claim 1 in prevention of graft-versus-host disease associated with allogeneic stem cell transplantation in a patient, which comprises the steps of:

- a) activating lymphocytes from a donor by mixing donor cells with host cells for a period of time sufficient for an immune reaction to occur;
- b) substantially eliminating the activated lymphocytes of step a) with photodynamic

DEC. 20. 2001 3:44PM SWABEY OGILVY MTL 514 288 8389

NO. 9406 P. 13/17

- 44 -

therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and

c) performing allogenic stem cell transplantation using the treated mix of step b).

11. Use of a composition according to claim 1 the treatment of immunologic disorder in a patient, which comprises the steps of:

- a) harvesting said patient's hematopoietic cells;
- b) ex vivo treating of the hematopoietic cells of step a) by photodynamic therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and
- c) performing graft infusion or autograft transplantation using the treated hematopoietic cells of step b).

12 Use according to claim 11, wherein said immunologic disorder is selected from the group consisting of conditions in which self cells or donor cells react against host tissues or foreign targets, such as graft-versus-host disease, graft rejection, autoimmune disorders and immunoallergic conditions.

13. Use of a composition according to claim 1 for the manufacture of a medicament for the photodynamic treatment for the selective destruction and/or inactivation of immunologically reactive cells without substantially affecting the normal cells or causing systemic toxicity for the patient, wherein appropriate



DEC. 20. 2001 3:45PM

SWABEY OGILVY MTL 514 288 8389

NO. 9406 P. 14/17

- 45 -

intracellular levels of said derivatives are achieved and irradiation of a suitable wavelength and intensity is applied.

14. Use of a composition according to claim 1 for the manufacture of a medicament for the prevention of graft-versus-host disease associated with allogeneic stem cell transplantation in a patient, which comprises the steps of:

- a) activating lymphocytes from a donor by mixing donor cells with host cells for a period of time sufficient for an immune reaction to occur;
- b) substantially eliminating the activated lymphocytes of step a) with photodynamic therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and
- c) performing allogeneic stem cell transplantation using the treated mix of step b).

15. Use of a composition according to claim 1 for the treatment of immunologic disorder in a patient, which comprises the steps of:

- a) harvesting said patient's hematopoietic cells;
- b) ex vivo treating of the hematopoietic cells of step a) by photodynamic therapy using a therapeutic amount of a photoactivable composition of claim 1 under irradiation of a suitable wavelength; and

$\frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$

DEC. 20. 2001 3:45PM

SWABEY OGILVY MTL 514 288 8389

NO. 9406 P. 15/17

- 46 -

16. Use according to claim 15, wherein said immunologic disorder is selected from the group consisting of conditions in which self cells or donor cells react against host tissues or foreign targets, such as graft-versus-host disease, graft rejection, autoimmune disorders and immunoallergic conditions.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 April 2001 (12.04.2001)

PCT


(10) International Publication Number  
**WO 01/24824 A1**

- (51) International Patent Classification: **A61K 41/00**
- (21) International Application Number: **PCT/CA00/01142**
- (22) International Filing Date: **3 October 2000 (03.10.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
**60/157,790**      **5 October 1999 (05.10.1999)**      **US**
- (71) Applicants (for all designated States except US): **UNIVERSITE DE MONTREAL [CA/CA]; 2900 Edouard-Montpetit, Montréal, Québec H3T 1J4 (CA). HOPITAL MAISONNEUVE-ROSEMONT [CA/CA]; 5415, boul. de l'Assomption, Montréal, Québec H1T 2M4 (CA).**
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ROY, Denis-Claude [CA/CA]; 2444 Prudential, Laval, Québec H7K 2C4 (CA). GUIMOND, Martin [CA/CA]; 7004 Paul Letondal St., Montréal, Québec H1E 5P2 (CA). MOLFINO, Nestor, A. [CA/CA]; 530 Victoria Avenue, Westmont, Québec H3Y 2R5 (CA).**
- (74) Agent: **SWABEY OGILVY RENAULT; 1981 McGill College Avenue, Suite 1600, Montréal, Québec H3A 2Y3 (CA).**
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
- *With international search report.*
  - *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*


(54) Title: **RHODAMINE DERIVATIVES FOR PHOTODYNAMIC DIAGNOSIS AND TREATMENT**

(57) Abstract: The present invention relates to the use of the photoactivable derivatives for the photodynamic treatment for the selective destruction and/or inactivation of immunologically reactive cells without affecting the normal cells and without causing systemic toxicity for the patient, wherein appropriate intracellular levels of said derivatives are achieved and irradiation of a suitable wavelength and intensity is applied.

**WO 01/24824 A1**




*Fig. 1A*




*Fig. 1B*

2/12



*Fig. 2*

3/12



*Fig. 3*

4/12



# RESTING LYMPHOCYTES

TIME (min) 30

50



# ACTIVATED LYMPHOCYTES





*Fig. 4A*

5/12



RESTING LYMPHOCYTES

TIME (min) 70

90



ACTIVATED LYMPHOCYTES




*Fig. 4B*




6/12

RESTING LYMPHOCYTES

TIME (min) 110





ACTIVATED LYMPHOCYTES



*Fig. 4c*

7/12

RESTING LYMPHOCYTES



ACTIVATED LYMPHOCYTES








Fig. 5

8/12




*Fig. 6*


9/12




10/12




*Fig. 8A*



*Fig. 8B*




*Fig. 8C*



*Fig. 9A*

- 1 - CONTROL, 0J/CM2
- 2 - CONTROL, 5J/CM2
- 3 - RH123 0J/CM2
- 4 - RH123 5J/CM2
- 5 - TH9402+RBBE, 0J/CM2
- 6 - TH9402+RBBE, 5J/CM2
- 7 - RBBE, 0J/CM2
- 8 - RBBE, 5J/CM2
- 9 - RBOE, 0J/CM2
- 10 - RBOE, 5J/CM2



*Fig. 9B*

# Declaration, Power Of Attorney and Petition

Page 1 of 3

WE (I) the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

RHODAMINE DERIVATIVES FOR PHOTODYNAMIC DIAGNOSIS AND TREATMENT

the specification of which

☐ is attached hereto.

☒ was filed on October 5, 1999 as  
Application Serial No. US 60/157,790  
and amended on \_\_\_\_\_.

☒ was filed as PCT international application  
Number PCT/CA00/01142  
on October 3, 2000,

and was amended under PCT Article 19  
on December 20, 2001 (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No



We (I) hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

_____ (Application Number)	_____ (Filing Date)
_____ (Application Number)	_____ (Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____

And we (I) hereby appoint: Norman F. Oblon, Reg. No. 24,618; Marvin J. Spivak, Reg. No. 24,913; C. Irvin McClelland, Reg. No. 21,124; Gregory J. Maier, Reg. No. 25,599; Arthur I. Neustadt, Reg. No. 24,854; Richard D. Kelly, Reg. No. 27,757; James D. Hamilton, Reg. No. 28,421; Eckhard H. Kuesters, Reg. No. 28,870; Robert T. Pous, Reg. No. 29,092; Charles L. Gholz, Reg. No. 26,395; Vincent J. Sunderdick, Reg. No. 29,004; William E. Beaumont, Reg. No. 30,996; Robert F. Gnuse, Reg. No. 27,295; Jean-Paul Lavalleye, Reg. No. 31,451; Stephen G. Baxter, Reg. No. 32,884; Robert W. Hahl, Reg. No. 33,893; Richard L. Treanor, Reg. No. 36,379; Steven P. Weihrouch, Reg. No. 32,829; John T. Goolkasian, Reg. No. 26,142; Richard L. Chinn, Reg. No. 34,305; Steven E. Lipman, Reg. No. 30,011; Carl E. Schlier, Reg. No. 34,426; James J. Kulbaski, Reg. No. 34,648; Richard A. Neifeld, Reg. No. 35,299; J. Derek Mason, Reg. No. 35,270; Surinder Sachar, Reg. No. 34,423; Christina M. Gadiano, Reg. No. 37,628; Jeffrey B. McIntyre, Reg. No. 36,867; Paul E. Rauch, Reg. No. 38,591; William T. Enos, Reg. No. 33,128; and Michael E. McCabe, Jr., Reg. No. 37,182; our (my) attorneys, with full powers of substitution and revocation, to prosecute this application and to transact all business in the Patent Office connected therewith; and we (I) hereby request that all correspondence regarding this application be sent to the firm of OBLON, SPIVAK, MCCLELLAND, MAIER & NEUSTADT, P.C., whose Post Office Address is: Fourth Floor, 1755 Jefferson Davis Highway, Arlington, Virginia 22202.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

100 Denis-Claude Roy

NAME OF FIRST SOLE INVENTOR

Signature of Inventor

Date

Residence: 2444, Prudential

Laval, Quebec *CHX*

CANADA H7K 2C4

Citizen of: \_\_\_\_\_

Post Office Address: \_\_\_\_\_


*2002/3/15*

200 Martin GUIMOND  
NAME OF SECOND JOINT INVENTOR

  
Signature of Inventor

2002/3/26  
Date

300 Nestor MOLFINO  
NAME OF THIRD JOINT INVENTOR

  
Signature of Inventor

Date

NAME OF FOURTH JOINT INVENTOR

Signature of Inventor

Date

NAME OF FIFTH JOINT INVENTOR

Signature of Inventor

Date

Residence: 1696 Quarry Trace

Columbus, Ohio 43204

U.S.A. OH

Citizen of:

Post Office Address:

Residence: 8817 Bells Mill Road

Potomac, Maryland 20854

U.S.A. MD

Citizen of:

Post Office Address:

Residence:

Citizen of:

Post Office Address:

Residence:

Citizen of:

Post Office Address: